

# A new synthetic route to 5,6,11,12-tetraarylethynyltetracenes

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

© 2018 The Royal Society of Chemistry. A new synthetic route to 5,6,11,12-tetrakis(arylethynyl)tetracenes,  $\pi$ -extended rubrenes, was developed via [4 + 2] cycloadditions of dialkynylisobenzofuran and 1,4-naphthoquinone. Introduction of arylethynyl groups by double nucleophilic additions to tetracenequinone gave sterically congested (arylethynyl)tetracenes after reductive aromatization. The photophysical properties of the newly prepared  $\pi$ -conjugated molecules are also evaluated.

<http://dx.doi.org/10.1039/c8ob02450b>

---

## References

- [1] K. Kitamura K. Asahina Y. Nagai H. Sugiyama H. Uekusa T. Hamura Chem.-Eur. J. 2018 24 14034 14038
- [2] T. Hamura, Patent JP 2016169213, 2016
- [3] For related reviews, see
- [4] M. M. Richter Chem. Rev. 2004 104 3003 3036
- [5] M. Bendikov F. Wudl D. F. Perepichka Chem. Rev. 2004 104 4891 4946
- [6] J. E. Anthony Angew. Chem. 2008 120 460 492 Angew. Chem., Int. Ed. 2008 47 452 483
- [7] For related distorted  $\pi$ -crowded ethynyl naphthalene derivatives, see
- [8] B. Bossenbroek D. C. Sanders H. M. Curry H. Shechter J. Am. Chem. Soc. 1969 91 371 379
- [9] H. A. Staab J. Ipaktschi Chem. Ber. 1971 104 1170 1181
- [10] A. E. Jung G. M. J. Schmidt Chem. Ber. 1971 104 3272 3288
- [11] See also
- [12] R. A. Pascal, Jr. Chem. Rev. 2006 106 4809 4819
- [13] T. Hamura R. Nakayama Chem. Lett. 2013 42 1013 1015
- [14] K. Asahina S. Matsuoka R. Nakayama T. Hamura Org. Biomol. Chem. 2014 12 9773 9776
- [15] R. Kudo K. Kitamura T. Hamura Chem. Lett. 2017 46 25 28
- [16] For our synthetic application of isobenzofurans to polyacene derivatives, see
- [17] S. Eda T. Hamura Molecules 2015 20 19449 19462
- [18] R. Akita K. Kawanishi T. Hamura Org. Lett. 2015 17 3094 3097
- [19] S. Eda F. Eguchi H. Haneda T. Hamura Chem. Commun. 2015 51 5963 5966
- [20] H. Haneda S. Eda M. Aratani T. Hamura Org. Lett. 2014 16 286 289
- [21] For syntheses of substituted rubrene derivatives, see
- [22] A. S. Paraskar A. R. Reddy A. Patra Y. H. Wijsboom O. Gidron L. J. W. Shimon G. Leitun M. Bendikov Chem.-Eur. J. 2008 14 10639 10647
- [23] J. Zhang S. Sarrafpour T. E. Haas P. Müller S. W. Thomas J. Mater. Chem. 2012 22 6182 6189
- [24] For a related [4 + 2] cycloaddition of diphenylisobenzofuran with 1,4-naphthoquinone, and structural characterization of the endo/exo-cycloadducts, see

- [25] T. Wombacher S. Foro J. J. Schneider Eur. J. Org. Chem. 2016 569 578
- [26] See also ref. 5 b and 6
- [27] The perfect endo selectivity observed in the [4 + 2] cycloaddition of isobenzofuran 3a and naphthoquinone 5 is presumably due to the lower solubility of the endo cycloadduct 6a in toluene than that of the exo cycloadduct 6a. Therefore, the endo cycloadduct 6a was selectively precipitated from the solution under the equilibrium between the starting materials and the cycloadducts. In a similar manner, the endo cycloadducts 6b-6d were obtained as a single stereoisomer by precipitation owing to their poor solubility with EtOH or heptane. For details, see ESI.
- [28] A similar approach for preparation of substituted rubrenes by using [4 + 2] cycloaddition of diarylisobenzofuran and naphthoquinone was developed, see
- [29] J. A. Dodge J. D. Bain A. R. Chamberlin J. Org. Chem. 1990 55 4190 4198
- [30] Under the acidic conditions, dialkynylisobenzofuran 3a was gradually decomposed, which promoted the cycloreversion of the cycloadduct 6a
- [31] For related base-induced aromatization, see
- [32] R. Polley M. Hanack J. Org. Chem. 1995 60 8278 8282
- [33] S.-H. Chan C.-Y. Yick H. N. C. Wong Tetrahedron 2002 58 9413 9422